

Hydrogen atom on curved noncommutative space

V.G. Kupriyanov*

CMCC, Universidade Federal do ABC, Santo André, SP, Brazil

September 28, 2012

Abstract

We have calculated the hydrogen atom spectrum on curved noncommutative space defined by the commutation relations $[\hat{x}^i, \hat{x}^j] = i\theta\hat{\omega}^{ij}(\hat{x})$, where θ is the parameter of noncommutativity. The external antisymmetric field which determines the noncommutativity is chosen as $\omega^{ij}(x) = \varepsilon^{ijk}x_k f(x_i x^i)$. In this case the rotational symmetry of the system is conserved, preserving the degeneracy of the energy spectrum. The contribution of the noncommutativity appears as a correction to the fine structure. The corresponding nonlocality is calculated: $\Delta x \Delta y \geq \frac{\theta^2}{4} m |\langle f^2 \rangle|$, where m is a magnetic quantum number.

1 Introduction

A combination of quantum mechanical arguments with general relativity indicates that at the distances of the order of the Planck length the space-time is nonlocal, therefore cannot be described as a differentiable manifold and should be treated as a kind of noncommutative structure [1]. Consider the noncommutative space realized by the coordinate operators $\hat{x}^i$, $i = 1, \dots, N$, satisfying the algebra

$$[\hat{x}^i, \hat{x}^j] = i\theta\hat{\omega}^{ij}(\hat{x}), \quad (1)$$

where $\hat{\omega}^{ij}(\hat{x})$ is an operator describing the noncommutativity of the space. The consistency condition for the algebra (1) implies, see e.g. [2], that the symbol $\omega_q^{ij}(x)$ of the operator $\hat{\omega}^{ij}(\hat{x})$ should have a form:

$$\omega_q^{ij}(x) = \omega^{ij}(x) + \omega_{co}^{ij}(x), \quad (2)$$

*e-mail: vladislav.kupriyanov@gmail.com

where $\omega^{ij}(x)$ is Poisson bi-vector, i.e., obey the Jacobi identity

$$\omega^{il}\partial_l\omega^{jk} + \omega^{kl}\partial_l\omega^{ij} + \omega^{jl}\partial_l\omega^{ki} = 0, \quad (3)$$

and the term $\omega_{co}^{ij}(x)$ stands for non-Poisson corrections to $\omega^{ij}(x)$ of higher order in θ , expressed in terms of $\omega^{ij}(x)$ and its derivatives, which depend on specific ordering of the operator $\hat{\omega}^{ij}(\hat{x})$. So, in order to determine the noncommutativity (1) we should define from physical considerations the antisymmetric field $\omega^{ij}(x)$ obeying the eq. (3) and specify the ordering. In what follows we treat $\omega^{ij}(x)$ as an external field and we choose the symmetric Weyl ordering.

This article aims to illustrate the possible phenomenological consequences of the presence of the noncommutativity of the type (1), when the commutator between coordinates is a function of these coordinates. As a starting point to study noncommutativity of the general form we consider quantum mechanics (QM). Two-dimensional model of position-dependent noncommutativity in QM was proposed in [3]. The particular example was considered in [4], where also was observed that canonical operators in these models are in general non Hermitian with respect to standard inner products [5]. For the example of QM and field theory on kappa-Minkowski space see e.g. [6] and references therein. The general form of QM with noncommutative coordinates of the type (1) was constructed in [7]. Note that the quantum mechanical scale of energies is rather different from the Planck scale, however some important properties like preservation of symmetries and corresponding consequences can be studied already in QM.

In particular, it is well known fact that the canonical noncommutativity, $[\hat{x}^i, \hat{x}^j] = i\theta^{ij}$, where θ^{ij} is an antisymmetric constant matrix, breaks the rotational symmetry of the Hydrogen atom, which removes the degeneracy of the energy levels [8]. This fact leads to the bounds of noncommutativity in this model. The same logic remains in the field theory, see e.g. [9] and references therein. We will show here that the noncommutativity can be introduced in a way to preserve the symmetries and the corresponding degeneracy. To this end in Section 2 we describe the model of QM with NC coordinates [7] and construct an explicit form of the trace functional on the algebra of the star product corresponding to (1). In Section 3 we consider a particle in a central potential on curved NC space, defined by the external antisymmetric field $\omega^{ij}(x) = \varepsilon^{ijk}x_k f(x_i x^i)$. For simplicity we set $(m_e = c = \hbar = 1)$.

2 Quantum mechanics with noncommutative coordinates

To define the QM on the noncommutative spaces of the type (1) we will need to introduce two objects, the star product and the trace functional. The existence of the star product

for any Poisson bi-vector $\omega^{ij}(x)$ is guaranteed by the Formality Theorem by Kontsevich [10], the explicit construction of the star product up to the fifth order in deformation parameter can be found in [2]. In particular, for any two functions f and g it has a form

$$(f \star g)(x) = f(\hat{x})g(x) = fg + \frac{i\theta}{2}\partial_i f \omega^{ij} \partial_j g - \frac{\theta^2}{4} \left[\frac{1}{2} \omega^{ij} \omega^{kl} \partial_i \partial_k f \partial_j \partial_l g - \frac{1}{3} \omega^{ij} \partial_j \omega^{kl} (\partial_i \partial_k f \partial_l g - \partial_k f \partial_i \partial_l g) \right] + O(\theta^3). \quad (4)$$

The trace functional is determined as $Tr(f) = \int \Omega(x) f(x)$, where $\Omega(x)$ is an integration measure. The trace should obey the property

$$Tr(f \star g) - Tr(fg) = 0, \quad (5)$$

which also guarantee the cyclic property of trace. The existence of a trace functional for a Kontsevich star-product related to any Poisson bi-vector ω^{ij} such that $div_\Omega \omega^{ij} = 0$ was demonstrated in [11]. But to make calculations one needs an explicit form for it. Suppose that there exist a function $\mu(x)$ such that

$$\partial_i (\mu \omega^{ij}) = 0. \quad (6)$$

Let $\Omega(x) = d^N x \mu(x)$, so

$$Tr(f) = \int d^N x \mu(x) f(x). \quad (7)$$

We should verify if this definition of trace is correct, i.e., if (5) holds true for any two functions f and g vanishing on the infinity. In the zero order in θ eq. (5) is just an identity, in the first order it is satisfied due to (6). In the second order in θ the right hand side of (5) is

$$-\frac{\theta^2}{4} \int d^N x \mu(x) \left[\frac{1}{2} \omega^{ij} \omega^{kl} \partial_i \partial_k f \partial_j \partial_l g - \frac{1}{3} \omega^{ij} \partial_j \omega^{kl} (\partial_i \partial_k f \partial_l g - \partial_k f \partial_i \partial_l g) \right] \quad (8)$$

Integrating this expression by parts on f and g and using (6) we rewrite it as

$$\frac{\theta^2}{12} \int d^N x \partial_i f \partial_l (\mu \omega^{ij} \partial_j \omega^{lk}) \partial_k g, \quad (9)$$

where the matrix $\partial_l (\mu \omega^{ij} \partial_j \omega^{lk})$ is symmetric, i.e.,

$$\partial_l (\mu \omega^{ij} \partial_j \omega^{lk}) = \partial_l (\mu \omega^{kj} \partial_j \omega^{li}), \quad (10)$$

due to the Jacobi identity and (6). The expression (9) is different from zero for two arbitrary functions f and g . To solve this problem we use the gauge freedom in the definition of the star product [10]. Let us construct a new star product

$$f \star' g = D^{-1} (Df \star Dg), \quad (11)$$

choosing a gauge operator D in such a way that (5) holds true for this new product. Taking into account (9) and (10) one should looking for D in the form

$$D = 1 + \theta^2 b^{ik} \partial_i \partial_k + O(\theta^3). \quad (12)$$

In this case

$$f \star' g = f \star g - 2\theta^2 b^{ik} \partial_i f \partial_k g + O(\theta^3). \quad (13)$$

The condition (5) for the star product (13) in the second order gives

$$\frac{\theta^2}{12} \partial_i f \partial_l (\mu \omega^{ij} \partial_j \omega^{lk}) \partial_k g - 2\theta^2 \mu b^{ik} \partial_i f \partial_k g = 0. \quad (14)$$

That is,

$$b^{ik} = \frac{1}{24\mu} \partial_l (\mu \omega^{ij} \partial_j \omega^{lk}). \quad (15)$$

We conclude that given a Poisson bi-vector ω^{ij} and a function $\mu(x)$ obeying (6), the modified star product (13) admits the trace (7).

Now, according to the [7] we describe the model of QM on NC spaces of general form, defined by the algebra (1). *The Hilbert space* is determined as a space of complex-valued functions which are square-integrable with a measure $\Omega(x)$. *The internal product* between two states $\varphi(x)$ and $\psi(x)$ from the Hilbert space is defined as

$$\langle \varphi | \psi \rangle = Tr(\varphi^* \star' \psi). \quad (16)$$

The action of the coordinate operators $\hat{x}^i$ on functions $\psi(x)$ from the Hilbert space is defined through the modified star product (13), for any function $V(x)$ one has

$$V(\hat{x}) \psi(x) = V(x) \star' \psi(x). \quad (17)$$

The definitions (16) and (17) means that the coordinate operators are self-adjoint with respect to the introduced scalar product: $\langle \hat{x}^i \varphi | \psi \rangle = \langle \varphi | \hat{x}^i \psi \rangle$. *The momentum operators $\hat{p}_i$* are fixed from the condition that they also should be self-adjoint with respect to (16). One of the possibilities is to choose it in the form

$$\hat{p}_i = -i\partial_i - i\partial_i \ln \mu(x). \quad (18)$$

One can easily verify that in this case $\langle \hat{p}_i \varphi | \psi \rangle = \langle \varphi | \hat{p}_i \psi \rangle$.

The momentum operators (18) commute, $[\hat{p}_i, \hat{p}_j] = 0$. The commutator between $\hat{x}^i$ and $\hat{p}_j$ is

$$[\hat{x}^i, \hat{p}_j] = i\delta_j^i - \frac{i\theta}{2} (\partial_j \omega^{il}(\hat{x}) \hat{p}_l + i\partial_j (\omega^{il} \partial_l \ln \mu)(\hat{x})) + O(\theta^2). \quad (19)$$

So, the complete algebra of commutation relations involving $\hat{x}^i$ and $\hat{p}_j$ is a deformation in θ of a standard Heisenberg algebra.

3 Particle in a central potential

Let us consider a particle placed in a central potential $V(r^2/2)$, where $r^2 = x^2 + y^2 + z^2$. The corresponding Hamiltonian reads

$$\hat{H} = \frac{\hat{p}^2}{2} + V\left(\frac{\hat{r}^2}{2}\right). \quad (20)$$

As it was already mentioned in the introduction, the canonical noncommutativity which can be realized by the coordinate operators $\hat{x}_{can}^i = x^i + i/2\theta^{ij}\partial_j$ violates the rotational symmetry of (20), since $[L_i, V(\hat{x}_{can}^2/2)] \neq 0$, where $L^i = -i\varepsilon^{ijk}x_j\partial_k$ is the angular momentum operator. And this problem cannot be solved introducing a noncommutativity as a Drinfeld twist deformation [12]. See also [13] for an example of nonabelian twist in QM.

The external antisymmetric field $\omega^{ij}(x)$ can be chosen in a way to preserve a rotational symmetry of the system. Let us suppose that

$$\omega^{ij} = \varepsilon^{ijk}x^k f(r^2), \quad (21)$$

where f is a given function. The corresponding noncommutative algebra is

$$[\hat{x}^i, \hat{x}^j] = i\theta\varepsilon^{ijk}\hat{x}^k f(\hat{r}^2). \quad (22)$$

The consistency condition for this algebra, $[\hat{x}^i, \varepsilon^{jkl}\hat{x}^l f(\hat{r}^2)] + cycl.(ijk) = 0$, is satisfied automatically and no corrections $\omega_{co}^{ij}(x)$ are needed to the (21) to construct the explicit form of the star product in any order in θ using the iterative procedure [2].

One can easily see that any function $\mu(r^2)$ obeys the equation (6):

$$\partial_i (\mu(r^2)\varepsilon^{ijk}x^k f(r^2)) = 0,$$

and can be chosen as a measure to define a trace functional. For simplicity we set $\mu(x) = 1$. So, the momentum operators are just derivatives $\hat{p}_i = -i\partial_i$.

The Hamiltonian (20) on the noncommutative space (22) will takes the form:

$$\begin{aligned}\hat{H} &= -\frac{1}{2}\Delta + V\star' = \\ &= -\frac{1}{2}\Delta + V - \frac{\theta^2}{12}V'f^2(r^2\Delta - 2x^i\partial_i - x^ix^j\partial_i\partial_j) + O(\theta^4) = \\ &= -\frac{1}{2}\Delta + V + \frac{\theta^2}{12}V'f^2L^2 + O(\theta^4),\end{aligned}\tag{23}$$

where $L^2 = L_x^2 + L_y^2 + L_z^2$ is the the orbital angular momentum. The specific choice of the external field (21) implies the conservation of the rotational symmetry:

$$\left[L_i, V\left(\frac{1}{2}\hat{r}^2\right)\right] = 0.\tag{24}$$

This, in turn, means the preservation of the degeneracy of the corresponding energy spectrum over the magnetic quantum number m .

In particular, for the Coulomb potential the leading order perturbation due to non-commutativity is

$$\hat{V}_{NC} = -\frac{e^2\theta^2}{12}\frac{f^2L^2}{r^3}.\tag{25}$$

We use the usual perturbation theory to calculate the leading corrections to the energy levels,

$$\Delta E_n^{NC} = \left\langle \psi^0 \left| \hat{V}_{NC} \right| \psi^0 \right\rangle.\tag{26}$$

where $|\psi^0\rangle = R_{nl}(r)Y_l^m(\vartheta, \varphi)$ is the unperturbed wave function, corresponding to the energy $E_n = -e^2/2a_0n^2$, here n is a principal quantum number and $a_0 = 1/\alpha$ is the Bohr radius. Integrating over the angular variables ϑ and φ , we stay with

$$\Delta E_n^{NC} = -\frac{e^2\theta^2l(l+1)}{12}\left\langle \frac{f^2}{r^3} \right\rangle,\tag{27}$$

where l is the azimuthal quantum number.

In order to calculate the corresponding nonlocality we first write (22) as

$$[\hat{x}^i, \hat{x}^j] = i\theta f \varepsilon^{ijk} x^k + \frac{i\theta^2}{2} f^2 \varepsilon^{ijk} L_k + O(\theta^3).\tag{28}$$

Taking into account that $\langle \psi^0 | f(r^2) x^k | \psi^0 \rangle = 0$, due to the integration over the angular variables, we end up with

$$\Delta x \Delta y \geq \frac{\theta^2}{4} m |\langle f^2 \rangle|,\tag{29}$$

where m is a magnetic quantum number.

We will consider two possibilities for the function f here: $f = 1$ and $f = r$. In the both cases the external field is zero at the origin and then grows along the corresponding axis, however in the second case the growth is higher. In the first case, taking into account the fact that

$$\left\langle \frac{1}{r^3} \right\rangle = \frac{1}{n^3 a_0^3} \frac{1}{l(l+1/2)(l+1)},$$

the the energy level shift is

$$\Delta E_n^{NC} = -\frac{\theta^2 E_n^2}{3a_0 e^2} \frac{n}{l+1/2}. \quad (30)$$

This energy correction has the same form as a relativistic kinetic energy contribution to the fine structure. Nonlocality in this case is defined by the relation

$$\Delta x \Delta y \geq \frac{\theta^2}{4} m. \quad (31)$$

Reminding that $\langle \frac{1}{r} \rangle = 1/n^2 a_0$, the energy correction for the second possibility ($f = r$) is given by

$$\Delta E_n^{NC} = \frac{\theta^2}{6} E_n l(l+1). \quad (32)$$

Finally since,

$$\langle r^2 \rangle = \frac{n^2 a_0^2}{4} (4n^2 - l^2 + 2nl + 1),$$

the corresponding nonlocality is given by

$$\Delta x \Delta y \geq m \frac{\theta^2 n^2 a_0^2}{16} (4n^2 - l^2 + 2nl + 1). \quad (33)$$

We can see that in this case the nonlocality depends on the energy of the system, the more the energy, the more the nonlocality.

4 Conclusions and perspectives

This example shows that the noncommutativity can be introduced in a minimal way in the theory, i.e., one may obtain nonlocality without violating physical observables like the energy spectrum, etc.

The considered model also indicates the way how to construct a relativistic generalization of QM with NC coordinates [7]. If the external field $\omega^{\rho\sigma}(x)$ which determine the

commutation relations $[\hat{x}^\rho, \hat{x}^\sigma] = i\theta\hat{\omega}^{\rho\sigma}(\hat{x})$, transforms as a two tensor with respect to a Lorentz group, the operators $\hat{x}^\rho = x^\rho + i\theta/2\omega^{\rho\sigma}\partial_\sigma + O(\theta^2)$ and $\hat{p}_\rho = -i\partial_\rho - i\partial_\rho\mu(x)$ will transform as vectors. This fact may be used to construct relativistic wave equations on curved NC space-time. For simplicity one may start with $(2+1)$ dimensions, choosing the external antisymmetric field as $\omega^{\rho\sigma}(x) = \varepsilon^{\rho\sigma\lambda}x_\lambda f(x^2)$.

Alternative models of relativistic QM with noncommutative coordinates were constructed in [14] in $(3+1)$ dimensions and in [15] in $(2+1)$ dimensions as a relativistic version of spin noncommutativity [16]. However, in this case the commutator between coordinates is proportional to the spin, i.e., coordinates do not form an algebra. This lead to the difficulties in the definition of the star product and makes the operators to be non Hermitian [17].

References

- [1] S. Doplicher, K. Fredenhagen and J. Roberts, Commum.Math.Phys. **172** (1995) 187.
- [2] V.G. Kupriyanov, D.V. Vassilevich, Eur.Phys.J.C. **58** (2008) 627-637.
- [3] M. Gomes, V.G. Kupriyanov, Phys.Rev.**D79** (2009) 125011.
- [4] A. Fring, L. Gouba, F.G. Scholtz, J.Phys.A A43 (2010) 345401.
- [5] B. Bagchi, A. Fring, Phys. Lett. A373 (2009) 4307.
- [6] E.Hariukumar, T.Juric, S. Meljanac, Phys.Rev.**D84** (2011) 085020; Phys.Rev.**D86** (2012) 045002.
- [7] V.G. Kupriyanov, *General form of quantum mechanics with noncommutative coordinates*, arXiv:1204.4823 [math-ph].
- [8] M. Chaichian, M.M. Sheikh-Jabbari and A. Tureanu, Phys.Rev.Lett. **86** (2001) 2716.
- [9] T.C. Adorno, D.M. Gitman, A.E. Shabad, D.V. Vassilevich, Phys.Rev.**D84** (2011) 085031.
- [10] M. Kontsevich, Lett. Math. Phys. **66** (2003) 157.
- [11] G. Felder, B. Shoikhet, Lett. Math. Phys. **53** (2000) 75 .
- [12] B. Chakraborty, Z. Kuznetsova, F. Toppan, J.Math.Phys. 51 (2010) 112102.
- [13] P.G. Castro, R. Kullock, F. Toppan, J.Math.Phys. 52 (2011) 062105.

- [14] M. Gomes, V.G. Kupriyanov, A.J.Silva, Phys.Rev.**D81** (2010) .
- [15] H.Falomir, F.Vega, J.Gamboa, F.Mendez, M.Loewe, *Noncommutativity in (2+1)-dimensions and the Lorentz group*, arXiv:1208.6315 [hep-th].
- [16] H.Falomir, J.Gamboa, J.Lopez-Sarrion, F.Mendez, P.Pisani, Phys.Lett **B680** (2009) 384.
- [17] A.F.Ferrari, M. Gomes, V.G. Kupriyanov, C.Shtehman, *Dynamics of a Dirac fermion in the presence of spin noncommutativity*, arXiv:1207.0412 [hep-th].